
22

## **CLAIMS**

## 1. A compound of formula I



wherein

each of X, Y, Z is independently selected from O, S, NR<sup>1</sup>, CR<sup>2</sup>R<sup>3</sup>, N and CR<sup>4</sup>, and where optionally X-Y, Y-Z, Z-E<sup>1</sup> and X-E<sup>2</sup> each independently form part of a saturated or unsaturated ring system which may be substituted or unsubstituted;

m is 0 or 1;

M is a metal selected from Ti[III], Ti[IV], Fe[II], Fe[III], Co[II], Co[III], Co[III], Ni[III], Cr[IIII], Mn[III], Mn[III], Mn[III], Ru[III], Ru[III], Ru[III], Pd[III], V[III], V[III], V[IVI], V[V], Cu[I], Cu[II], Rh[III], Rh[III], Mo[III], Mo[V], Re[I] and Re[III]; each of E<sup>1</sup> and E<sup>2</sup> is independently selected from O, S, NR<sup>5</sup>, N, P, PR<sup>6</sup>, where at least one of either E<sup>1</sup> or E<sup>2</sup> carries a formal negative charge;

L<sup>2</sup> is a one electron donor ligand;

n is zero or an integer such that the compound has an overall charge of zero or +1; L<sup>1</sup> is NR<sup>7</sup>R<sup>8</sup>, PR<sup>7</sup>R<sup>8</sup>, OR<sup>7</sup>, SR<sup>7</sup>, O, S or NR<sup>16</sup>, imidazolyl, pyridinyl, benzimidazolyl or quinolinyl;

each of R<sup>1-8</sup> and R<sup>16</sup> is independently H or a hydrocarbyl group;

Q is a linker group; and

r is 0 or 1.

- 2. A compound according to claim 1 wherein  $L^2$  is selected from halide, hydride, alkyl and cyanide.
- 3. A compound according to claim 1 or claim 2 wherein L<sup>2</sup> is chloride or bromide.

- 4. A compound according to any preceding claim wherein X, Y and Z are each independently selected from  $CR^2R^3$  and  $CR^4$ .
- 5. A compound according to claim 4 wherein:
- (i) m is 1, each of X-E<sup>2</sup> and Y-Z is independently a single or a double bond or part of a delocalised  $\pi$  system, and X-Y and Z-E<sup>1</sup> are single bonds; or
- (ii) m is 1, each of X-Y and Z-E<sup>1</sup> is independently a single or a double bond or part of a delocalised  $\pi$  system, and Z-E<sup>2</sup> and Y-Z are single bonds; or
- (iii) m is 0, each of  $X-E^2$  and  $Z-E^1$  is independently a single or a double bond or part of a delocalised  $\pi$  system, and X-Z is a single bond;
- 6. A compound according to any preceding claim wherein m is one, Y-Z is a double bond or part of a delocalised  $\pi$  system, and X-E<sup>2</sup> is a single or a double bond.
- 7. A compound according to any one of claims 1 to 6 which comprises a compound of formula II

$$R^{11}$$
 $R^{10}$ 
 $R^{10}$ 

wherein each of  $R^{9-14}$  is independently H, a hydrocarbyl group, a halide, ether, thioether, ester, nitro, dialkylamino, or cyano group, and "a" is a double bond or part of a delocalised  $\pi$  system (where one of  $R^9$  or  $R^{10}$  is absent), or "a" is a single bond.

8. A compound according to any preceding claim wherein  $X-E^2$  is a double bond or part of a delocalised  $\pi$  system, and  $E^2$  is N.

ï 24

- 9. A compound according to any one of claims 1 to 7 wherein X-E<sup>2</sup> is single bond and E<sup>2</sup> is NR<sup>5</sup>.
- 10. A compound according to any preceding claim wherein E<sup>1</sup> is O.
- 11. A compound according to any preceding claim which comprises a compound of formula III or IV

wherein each of R<sup>9-14</sup> is independently H, a hydrocarbyl group, a halide, ether, thioether, ester, nitro, amino, or cyano group.

- 12. A compound according to any preceding claim wherein M is Fe.
- 13. A compound according to any preceding claim wherein  $L^2$  is chloride and n is one or two.
- 14. A compound according to any one of claims 1-5, 8-10 or 12-13 wherein m is 0,  $X-E^2$  and  $Z-E^1$  are both double bonds or each form part of a delocalised  $\pi$  system, and X-Z is a single bond.
- 15. A compound according to claim 14 wherein said compound is of formula V, VI or VII

wherein each of  $R^{9-14}$  is independently H, a hydrocarbyl group, a halide, ether, thioether, ester, nitro, dialkylamino, or a cyano group

16. A compound according to any preceding claim wherein  $L^1$  is selected from the following: O, -S, -NR<sup>16</sup>,

$$\stackrel{\downarrow}{\leftarrow} \text{NMe}_2$$
 $\stackrel{\downarrow}{\leftarrow} \text{NMe}_2$ 
 $\stackrel{\downarrow}{\leftarrow} \text{SPh}$ 
 $\stackrel{\downarrow}{\leftarrow} \text{OPh}$ 
 $\stackrel{\downarrow}{\leftarrow} \text{OMe}$ 
 $\stackrel{\downarrow}{\leftarrow} \text{PPh}_2$ 

- 17. A compound according to any preceding claim wherein the linker group Q is  $(CHR^{15})_{p}$  or a phenylene group, where p is 1, 2, 3....10, and each  $R^{15}$  is independently H or a hydrocarbyl group.
- 18. A compound according to claim 17 wherein the linker group Q is o-phenylene or -(CH<sub>2</sub>)<sub>p</sub>- where p is 1 or 2.
- 19. A compound according to any preceding claim wherein r is 1.
- 20. A compound according to any preceding claim wherein each of  $R^{1-15}$  is independently a  $C_{1-50}$  alkyl optionally comprising one or more heteroatoms, aryl, or a heteroaryl.

- 21. A compound according to any preceding claim wherein each of  $R^{1-15}$  is independently a  $C_{1-20}$  alkyl.
- 22. A compound according to any preceding claim wherein said compound of formula I is selected from the following:

- 23. A catalyst composition comprising a compound according to any one of claims 1 to 22 and an initiator.
- 24. A catalyst composition according to claim 23 wherein the initiator has a radically transferable atom or group.
- 25. A catalyst composition according to claim 23 or claim 24 wherein the initiator is selected from an alkyl halide optionally containing an electron withdrawing group in the alpha position, a substituted or unsubstituted arenesulphonyl halide, an alkyl dihalide, a sulphonyl halide and a polymer bearing one or more radically transferrable group
- 26. A catalyst composition according to claim 23 or claim 24 wherein the initiator is selected from CCl<sub>4</sub>, CHCl<sub>3</sub>, CCl<sub>3</sub>Br, 2-bromoethylisobutyrate, 2-

bromoisobutyrophenone, para-toluenesulphonyl chloride, phenoxybenzene-4,4'-disulphonyl chloride, 1,3-benzene disulphonyl chloride.

- 27. A catalyst composition according to claim 23 or claim 24 wherein the initiator is AIBN.
- 28. A catalyst composition according to any one of claims 23 to 27 wherein the compound of formula I is supported on an inorganic or organic solid support.
- 29. Use of a compound according to any one of claims 1 to 22, or a catalyst composition according to any one of claims 23 to 28, for polymerising a radically polymerisable monomer.
- 30. A process for polymerising a radically polymerisable monomer, said process comprising contacting a catalyst composition according to any one of claims 23 to 28 with a radically polymerisable monomer, optionally in the presence of a solvent.
- 31. A process according to claim 30 wherein the radically polymerisable monomer is selected from one or more of the following:  $C_{2-8}$  alpha olefins, optionally substituted conjugated dienes, acrylic acid, acrylic anhydride, (meth)acrylamides, vinyl halides, (meth)acrylonitrile, (meth)acrylate esters of  $C_{1-20}$  alcohols, vinyl esters of  $C_{1-20}$  alcohols, vinyl amides having up to 8 carbons, vinyl ketones having up to 8 carbons, vinyl substituted aryls.
- 32. A process according to claim 30 or claim 31 wherein the radically polymerisable monomer is an acrylate selected from the following: methyl acrylate, ethyl acrylate, butyl acrylate, 2- ethylhexyl acrylate, isobornyl acrylate, and functional derivatives thereof such as 2- hydroxy ethyl acrylate, 2-chloro ethyl acrylate.
- 33. A process according to claim 30 or claim 31 wherein the radically polymerisable monomer is a methacrylate selected from the following: methyl methacrylate, ethyl methacrylate, butyl methacrylate, 2-ethylhexyl methacrylate,

isobornyl methacrylate, 2-hydroxy ethyl methacrylate, 2-chloro ethyl methacrylate, 2-hydroxypropyl methacrylate, (HPMA) 2-morpholinoethylmethacrylate (MEMA), 2-(dimethylamino)ethyl methacrylate (DMA), glyerol monomethacrylate (GMA), methoxy capped oligo(ethyleneglycol) methacrylate (OEGMA), poly(ethyleneglycol) methacrylate (PEGMA), glycidyl methacrylate.

- 34. A process according to claim 30 or claim 31 wherein wherein the radically polymerisable monomer is a (meth)acrylamide selected from the following: (meth)acrylamide, N-methyl (meth)acrylamide and, N,N'dimethyl (meth)acrylamide.
- 35. A process according to claim 30 or claim 31 wherein the radically polymerisable monomer is selected from the following: styrene, methyl acrylate, methyl methacrylate, 2-hydroxypropyl methacrylate, (HPMA) 2-morpholinoethylmethacrylate (MEMA), 2-(dimethylamino)ethyl methacrylate (DMA), glyerol monomethacrylate (GMA), methoxy capped oligo(ethyleneglycol) methacrylate (OEGMA), poly(ethyleneglycol) methacrylate (PEGMA) and glycidyl methacrylate.
- 36. A process according to any one of claims 30 to 35 wherein the ratio of initiator to radically polymerisable monomer is from  $2\times10^{-3}$ :1 to  $1\times10^{-4}$ :1.
- 37. A process according to any one of claims 30 to 35 wherein the ratio of initiator to radically polymerisable monomer is from  $1\times10^{-3}$ :1 to  $1.6\times10^{-4}$ :1.
- 38. A process according to any one of claims 30 to 35 wherein the ratio of initiator to radically polymerisable monomer is from  $4\times10^{-4}$ :1 to  $2\times10^{-4}$ :1.
- 39. A process according to any one of claims 30 to 38 wherein the ratio of initiator to the compound of formula I is from  $1 \times 10^{-4}$ :1 to 10:1.
- 40. A process according to any one of claims 30 to 39 wherein the ratio of initiator to the compound of formula I is from  $1 \times 10^{-1}$ :1 to 5:1.

- 41. A process according to any one of claims 30 to 40 wherein the polymerisation takes place at a temperature of from about -20°C to 200°C.
- 42. A process according to any one of claims 30 to 41 wherein the polymerisation takes place in the presence of a Lewis acid activator.
- 43. A process according to claim 42 wherein the Lewis acid activator is an aluminium alkyl, an aluminium alkoxide, an aluminium halide an alkyl zinc reagent, or a borane.
- 44. A process according to claim 43 wherein the Lewis acid activator is selected from methyl aluminium, bis(2,6 di-tert-butylphenoxide), aluminium tris(iso-propoxide), aluminium trichloride, diethyl zinc, BPh<sub>3</sub> and B( $C_6F_5$ )<sub>3</sub>.
- 45. A process according to any one of claims 42 to 44 wherein the ratio of activator to the compound of formula I is from 1:1 to 10:1.
- 46. A process according to any one of claims 29 to 44 wherein the polymerisation is carried out in bulk, solution, emulsion, suspension or in the gas phase.
- 47. A polymerisation mixture comprising a catalyst composition according to any one of claims 23 to 28 and a radically polymerisable monomer as defined in any one of claims 31 to 35, which optionally further comprises a solvent and/or a Lewis acid activator.